Rhodora

JOURNAL OF THE

NEW ENGLAND BOTANICAL CLUB.

Conducted and published for the Club, by

BENJAMIN LINCOLN ROBINSON, Editor-in-chief.

FRANK SHIPLEY COLLINS
MERRITT LYNDON FERNALD
HOLLIS WEBSTER } Associate Editors.

WILLIAM PENN RICH
EDWARD LOTHROP RAND } Publication Committee.

Vol. 15. November, 1913. No. 179.

CONTENTS

The Heather on Martha's Vineyard. *E. P. Bicknell* . . 189

Two new Species of Stigonema. *F. N. Blanchard* . . . 192

Six Weeks' Botanizing in Vermont,— II. *S. F. Blake* . . 200

A new Station for Scirpus Longii. *M. L. Fernald* . . . 202

The Josselyn Botanical Society. *D. W. Fellows* . . . 202

Field Meeting of the Vermont Botanical Club. *N. F. Flynn* . 203

A Correction concerning Dicksonia. *M. L. Loomis* . . . 204

Boston, Mass. 1052 Exchange Building.

Providence, R. I. Preston and Rounds Co.

RHODORA.—A monthly journal of botany, devoted primarily to the flora of New England. Price ***$1.50*** per year (domestic and foreign); single copies 15 cents. Prices of Volumes 1 and 2 (1899 and 1900) on application. All remittances by check or draft, except on Boston or New York, must include ten cents additional for cost of collection. Notes and short scientific papers, relating directly or indirectly to the plants of the northeastern states, will be gladly received and published to the extent that the limited space of the journal permits. Forms will be closed five weeks in advance of publication. Authors (of more than one page of print) will receive 25 copies of the issue in which their contributions appear. Extracted reprints, if ordered in advance, will be furnished at cost.

Address manuscripts and proofs to
B. L. ROBINSON, 3 Clement Circle, Cambridge, Mass.

Subscriptions, advertisements, and business communications to
W. P. RICH, 300 Massachusetts Avenue, Boston, Mass.

Single copies may be had from
E. L. RAND, Corresponding Sec'y N. E. Botanical Club,
1052 Exchange Building, Boston, Mass.

Entered at Boston, Mass., Post office as Second Class Mail Matter

---

KEY TO NEW ENGLAND TREES, Wild and Commonly Cultivated, based primarily upon leaf characters, by J. FRANKLIN COLLINS and HOWARD W. PRESTON. Price 40c. net. PRESTON & ROUNDS CO., Providence, R. I.

CARD-INDEX OF NEW GENERA, SPECIES AND VARIETIES OF AMERICAN PLANTS, 1885 TO DATE.

For American taxonomists and all students of American plants the most important supplement to the Index Kewensis, this catalogue in several ways exceeds the latter work in detail, since it lists not merely the flowering plants, but pteridophytes and cellular cryptogams, and includes not merely genera and species, but likewise subspecies, varieties and forms. A work of reference invaluable for larger herbaria, leading libraries, academies of sciences, and other centers of botanical activity. Issued quarterly, at $15.00 per 1000 cards.

GRAY HERBARIUM of Harvard University,
Cambridge, Mass., U. S. A.

CHECK LIST OF GRAY'S MANUAL, 7th EDITION, compiled by M. A. DAY. Leatherette. Pocket size. Invaluable for collectors' memoranda and herbarium records. Published and sold by the GRAY HERBARIUM, Cambridge, Mass. Price postpaid 20 cts. each. Ten copies $1.50.

Advertisements of Nurserymen and Dealers in Botanical and other Scientific Publications are inserted in these pages at the following rates per space of 4 in. by 3–4 in. 1 year $4.00. 6 months $2.50.

Rhodora

JOURNAL OF

THE NEW ENGLAND BOTANICAL CLUB

Vol. 15. November, 1913. No. 179.

THE HEATHER, CALLUNA VULGARIS, ON MARTHA'S VINEYARD.

Eugene P. Bicknell.

So much has been made of the heather growing wild on Nantucket that a sort of distinction has come to that island by reason of this plant. As well might the plural form be used — of these plants — for, in addition to the better known common Heather or ling (*Calluna vulgaris* (L.) Salisb.) two other heaths are found on Nantucket — the Bell Heather (*Erica cinerea* L.) and the Cross-leaved Heath (*Erica tetralix* L.). In respect of the *Calluna* it is to be reported that the island of Martha's Vineyard has equal if not greater claim to honor. Yet so little had this been suspected that when I chanced upon the heather growing there, in far greater profusion and in altogether wilder surroundings than on Nantucket, it seemed by first impression almost like an intrusion on a prerogative of the more seaward island.

Among the botanical elect of Nantucket its heather secrets have long been held in loyal reserve. But for this circumspection it is doubtful if any heather at all would be growing wild there today. It cannot be thought ungracious therefore if I particularize here no more fully as to the Martha's Vineyard locality than to report it as being in pitch pine woods in the southeastern quarter of the island.

An obscure woodland road passes the spot not many rods distant and an abandoned farm house, not yet a ruin, occupies an old clearing not very far away, but the general surroundings are quite uninhabited tracts of scrub oak, and woodland of oak and pine, traversed by sandy cartways and remote from any highway or any cultivated ground. The main growth of the heather carpets the level floor of

the woods over an irregular area some twenty-seven paces in length and twenty-four paces in greatest breadth. No garden spot could show a more perfectly continuous bed of color than here lies hidden among the pines in September when the heather is in full bloom. On three sides it is well screened by thick woodland growth, but from one direction no alert botanist in passing would fail to catch glimpses of its rosy glow through openings among the trees while yet eight or ten rods away. Of the few outlying patches the most distant is about fifty paces from the favored spot; another, thirty-five paces distant, lies quite outside the grove-like tract of pines in the adjoining scrub oak. In September, 1912, it was found that this patch had been discovered and much of it had been torn up and left lying about to die. The main growth, still undisturbed, had evidently been established for many years. The larger pines that gave it partial shade were from twenty to nearly thirty feet in height. When discovered, on September 24, 1909, it was in full flower; on October 10 fresh flowers were still appearing but the mass of bloom had lost its bright color of two weeks before. Few other kinds of plants grew close about it; the most noteworthy were *Cypripedium acaule*, *Hypopitys lanuginosa*, *Monotropa uniflora*, *Vaccinium pennsylvanicum*, *Epigaea repens*, *Pyrola rotundifolia*, *Trientalis americana* and *Carex pennsylvanica*.

This Martha's Vineyard heather appeared so different from the Nantucket plant — more straggling in habit with shorter less tapering racemes of deeper pink and somewhat different flowers and, in effect, glabrous foliage — that it gave me the instant impression of being quite like a different species. Nor was this impression an altogether misleading one. The plant is indeed true *Calluna vulgaris*, that is to say, the typical glabrate form of the species. The Nantucket *Calluna*, on the other hand, proves to be the markedly pubescent form of the plant which by some European botanists has been esteemed a distinct variety or even a valid species. It is the *Calluna vulgaris* var. *pubescens* of Koch (Syn. ed. 2: 547, 1844).

It does not clearly appear from the specimens examined how worthy of distinction from the more glabrous type this pubescent heather may be, or whether the minute pubescence of its younger parts and sometimes even of the leaves, is to be taken as evidence that the two plants readily interblend. The glabrate plant appears to be the prevailing and more widely diffused form in Europe. In the collections at the New York Botanical Garden I find specimens from England, France,

Germany, Hungary and Iceland, the pubescent form appearing from England only. But it is to be noted that it is reported by Schur from Kronstadt (Enum. Pl. Transs. 447, 1866).—"*Calluna vulgaris* b. *pubescens* Koch. (*Erica ciliaris* Hudson [non L.]—*Calluna ciliaris* Schur, herb. Transs.)." Specimens of the glabrate plant from America in the same Herbaria are as follows: Massachusetts — Andover, 1861, ex herb. A. Gray; Tewksbury, 1874, Thos. Morong; Nantucket; Rhode Island — Worden's Pond, Sept., 1894, J. F. Kemp; New Jersey — near Hammondton, Aug., 1891, W. A. Stowell. That the glabrate plant is found also in Newfoundland is clearly shown by the figure in Journ. Bot. **4**: 306, 1866, where the American heather was mistakenly proposed as distinct from the European under the name *Calluna atlantica* by Doctor Seemans.

The only notably pubescent specimens seen from any place in America are from Nantucket. There the plant has held its pubescent character from the time of its introduction thirty-six years ago. The evidence is interesting. There is preserved in the Herbarium of Columbia University an old Nantucket specimen, ex herb. O. R. Willis, bearing on the label no date or record other than the name of the collector, Mrs. C. C. Pearson. Mr. Willis seems to have been the first to have reported this heather, and also of *Erica Tetralix* from Nantucket. His note of announcement was published in the Bulletin of the Torrey Botanical Club for December, 1886, and records that the discovery of both species was made that same year by Mrs. Charlotte C. Pearson who had sent him specimens. This establishes beyond question the date and history of the Willis specimen. It is an excellent example of the pubescent form, and was from the same station where that form is found today.

It would seem to follow that no connection is to be supposed between the introduction of this pubescent heather on Nantucket and that of the glabrate form on Martha's Vineyard. There is even good reason for believing that the *Calluna* on Nantucket, even at the time of its earliest discovery there, must have come to the island from two different lines of approach, for both forms of the plant have been found there. A cluster that I came upon on the open plains in June, 1909, far away from the locality of the pubescent plant, was of the glabrate variety. Mrs. Owen has told us (RHODORA, **10**: 173–179, 1908) that it has been sought to spread the heather on Nantucket by scattering seed and even by setting out young plants. The existence of this isolated

cluster is perhaps thus to be explained. But there is evidence that the glabrate plant grew on Nantucket long before any effort had been made to introduce the heather there. The Herbarium of Columbia University contains another old specimen of *Calluna* labeled simply "Nantucket" without other record. It belongs among the earlier collections of the heather on that island and is of the glabrate form. It would seem to be most unlikely that both forms had come to the island together from the same place in Europe, and it is therefore to be inferred that Nantucket has received this addition to its flora from at least two sources of origin. Indeed Mrs. Owen believes (loc. cit.) that a solitary plant of *Calluna* found on Nantucket in 1880 far away from the locality where it was brought in with European conifers three years before was not of that introduction. How it came there is not less a mystery than is the presence of the heather at the other widely separated localities from Newfoundland to New Jersey where it has been found on the American Continent.

New York City.

TWO NEW SPECIES OF STIGONEMA.[1]

Frank N. Blanchard.

(Plate 105.)

In some material collected in October, 1909, by Dr. F. D. Lambert of Tufts College, from Chebacco Pond in the town of Essex, Essex County, Massachusetts, there was found very abundantly a blue-green alga, that apparently had not been described before, belonging to the family *Stigonemaceae*. This material was put into formalin and left until November, 1912, when Dr. Lambert and myself secured fresh material from the same place and found the same alga still plentiful. In April, 1913, I visited the pond and found the alga very scarce, but in exactly the same growing condition as in the previous November. It was found free-floating among other algae, chiefly blue-greens, where dead leaves and stems had collected in masses at the edge of the pond. Its filaments form loose, wiry-looking clusters from one to several millimeters in diameter.

[1] Contributions from the Biological Laboratories of Tufts College, No. 55.

The plant in the vegetative condition consists of blue-green filaments, 20–36 microns wide, that are repeatedly branched in every direction. Occasionally three or four branches may arise from adjacent cells, but the branches are usually more scattered. Although it is usually true that a branch is narrower than the filament it comes from, it may be equal to it in width, but never greater. In all the material collected the majority of branches appeared to be developed for the sole purpose of forming hormogones. No free vegetative ends of filaments, that were not forming or that had not already formed hormogones, were observed in all the material examined. The filaments are usually composed of a single row of cells, but two cells very frequently occur side by side. The colorless connections between the cells are plainly shown in formalin preserved material (fig. 8).

Lateral heterocysts are common (fig. 3). Intercalary heterocysts often occur just below a hormogone, and sometimes in other places, but they are less common than the lateral heterocysts. They contain no granules and vary in color from a light brown to a very dark blue.

The sheath is usually colorless, but many times is tinged at the ends of branches with a light, golden brown. The sheath varies from four to eight microns in thickness.

Filaments grow in length and develop branches by simple division of cells. Early stages in formation of a branch are shown in figures 1, 2, and 3. Hormogones are found of all lengths up to 196 microns, and their width is the same as that of the vegetative cells. They are developed in special branches and at the ends of main branches; they may occupy the whole of a branch and project down into the main filament, or may occupy only a part of the branch (figs. 3 and 6). When the plant is most actively forming hormogones, the vegetative cells become more and more vacuolated (figs. 1, 2, 3, and 6), and after the hormogones have escaped, the vegetative cells, sometimes if not always, degenerate. In one plant observed, every hormogone had escaped, and the sheath and cells had become a uniform brown, showing evident degeneration after escape of hormogones. Intermediate stages in this degeneration are easy to find.

The hormogones escape by a breaking off of the end of the sheath as in figure 6, or by apparent disorganization of the end of the sheath. The actual discharge of the hormogone is a slow process and appears to be accompanied by the discharge of a mucilaginous mass (figs. 4 and 5). The hormogone may escape from the sheath as a whole or

may break up into groups of one, two or more cells, either before or after leaving the sheath, as shown in figures 4, 5, and 7. In very young hormogones, the line of division between the cells is not clear, but as they grow older this line of separation increases in distinctness. The color of the hormogones changes as they mature, from the light blue-green of the vegetative cells to a very dark blue-green, which under the microscope appears almost black. This is probably due to a concentration of material. For this alga I propose the name

Stigonema **anomalum** sp. nov. Filis liberis, inter algas varias sparsis, repetite et irregulariter vel subsecundatim ramosis, 20–36 μ crassis; ramis patentibus, filo primario nunc aequicrassis, nunc tenuioribus, omnibus hormogoniferis; vagina 4–8 μ crassa, continua, hyalina vel ad apices aureo-fusca; cellulis diametro brevioribus, rectangularibus vel disciformibus, uni- vel biseriatis, aerugineis; heterocystis lateralibus, rarius intercalaribus, fuscis vel coeruleis; hormogoniis terminalibus, longitudine variis, usque ad 196 μ, aerugineis.

Filaments unattached, floating among other algae, repeatedly branched in every direction, or somewhat secundly, 20–36 μ diam.; branches patent, of the same size as the primary filament or sometimes thinner, all producing hormogones; sheath 4–8 μ thick, even, hyaline or sometimes golden brown at the apex of a branch; cells shorter than their diameter, mostly disciform or rectangular, of one or two series of cells, blue green; heterocysts lateral, less commonly intercalary, from brownish to blue-green; hormogonia terminal, of varying length, up to 196 μ.

Forming loose clumps, free floating among other, mostly blue-green, algae, Chebacco Pond, Essex, Massachusetts.

S. anomalum is a typical *Stigonema* for the following reasons: It is composed of one or two rows of cells; it is repeatedly branched in every direction; all branches bear hormogones; the cells are usually shorter than wide; the sheath is sometimes colored a light brown. It is allied to *Fischerella* and *Hapalosiphon* by its tendency to unilateral branching; by its smooth sheath; by the first cell of a branch projecting into the main filament; and by its generally rectangular cells.

Stigonema ocellatum is the only plant that could be confused with *S. anomalum*. I examined the specimen labeled *S. ocellatum* collected by G. T. Moore, in 1897 and distributed in Collins, Holden & Setchell, Phyc. Bor.-Am., No. 455, and concluded that my plant was not a new one after all, as it agreed very closely with Moore's plant. I then examined, through the kindness of Mr. Charles Bullard, the specimen of *S. ocellatum* from the Farlow Herbarium, originally from the herbarium of Bornet and Thuret, and decided that Moore had wrongly identi-

fied the plant he sent out, and that he had distributed *S. anomalum* under the name of *S. ocellatum*. *S. anomalum* has main filaments of about the size of *S. ocellatum;* as in that species, they consist usually of a single series of cells, occasionally two cells being side by side; otherwise, there is little resemblance. *S. anomalum* differs from *S. ocellatum* in the following ways:

1st. The wall is not very thick; the outline is straight, not following the individual cells.

2nd. There is no lamination either of the general wall, or about the individual cells. This latter, annular lamination, is what gives *S. ocellatum* its specific name.

3rd. The wall is usually colorless; in *S. ocellatum* it is generally colored, sometimes very strongly.

4th. In *S. anomalum* the cells are not rounded, but are mostly disciform.

5th. The branching is repeated, usually with diminution of diameter.

6th. The hormogones are generally longer than in *S. ocellatum.*

7th. The way the branches arise in *S. anomalum* is rather of the *Hapalosiphon* type, the one or two cells in the main filament appearing to belong to the branch rather than to the main filament.

Specimens of *Stigonema anomalum* will be distributed in the Phycotheca Boreali-Americana; the material was preserved in formalin before it was dried, and somewhat shrunken specimens are the result.

In the material collected in 1909, as above described under *Stigonema anomalum,* there was also found a new species of *Hapalosiphon.* Although there is no doubt that it is a *Hapalosiphon,* yet it is even more an intergrading form between *Hapalosiphon* and *Stigonema* than is *S. anomalum.* It was found free-floating under the same conditions as *S. anomalum,* but in appearance it is a small, soft sod a few millimeters across, of a blue-green to brownish color.

In the vegetative condition the plant consists of blue-green or brownish filaments from eight to sixteen microns wide that are repeatedly branched in every direction. There is a tendency for the branching to be unilateral as in *S. anomalum,* and for the secondary branches to be less in number than the primary branches. The branches do not taper toward the extremities, but are sometimes slightly clavate. The branches may or may not be slightly narrower than the primary filament. The cells are nearly always disposed in a single row. The

exception shown in figure 9 is rare. All the figures of this plant were made from material preserved in formalin and although there is very little shrinkage, yet there is just enough to show, in many places, the colorless connections between the cells. The cells are mostly cylindrical, usually longer than wide, and closely packed in the sheath. The colorless intercalary heterocysts are common (fig. 10). The sheath is mostly smooth, always colorless, and from two to four microns in thickness. At the ends of hormogone-bearing branches, the sheath is often thickened up to twelve microns (fig. 12). The cells of the branches are sometimes longer than those in the main filament, but are often shorter.

Filaments grow in length and develop branches by simple division of cells as shown in figure 9. Hormogones 37–163 microns in length and 8–12 microns in width occur at the ends of all branches or in short special branches. The cells of the hormogones are not well marked off from each other. Their color remains blue-green throughout their growth. The short hormogone-bearing branches are all variously curved (fig. 10).

Stigonema **medium** sp. nov. Filis liberis, inter algas varias sparsis, repetite et irregulariter vel subsecundatim ramosis, 8–16 μ crassis; ramis patentibus, filo primario plerumque tenuioribus, cylindricis vel clavatis; vagina 4–8 μ crassa, ad apices usque ad 12 μ, continua, hyalina; cellulis plerumque diametro longioribus, cylindricis, uniseriatis, aerugineis vel fuscis; heterocystis forma et magnitudine cellulis normalibus similibus, hyalinis, intercalaribus; hormogoniis terminalibus, 37–163 $\times$ 8–12 μ, aerugineis.

Filaments unattached, floating among other algae, repeatedly branched in all directions or somewhat secundly, 8–16 μ diam.; branches patent, generally thinner than the primary filament, cylindrical or clavate; sheath 4–8 μ thick, up to 12 μ at the ends of the branches, even, hyaline; cells usually longer than their diameter, cylindrical, in one series, blue-green or brownish; heterocysts similar to ordinary cells in form and size, hyaline, intercalary; hormogonia terminal, 37–163 $\times$ 8–12 μ, blue-green.

Forming a minute, soft sod, among other algae, mostly blue-green, Chebacco Pond, Essex, Massachusetts, October, 1909.

This plant shows characteristics of both *Hapalosiphon* and *Stigonema*. It is characteristically a *Hapalosiphon* from its single row of cylindrical cells usually longer than broad; from its method of branching, and in that it usually has the cells in the branches longer than those in the main filament. It has the characteristics of a *Stigonema* in its habit of forming hormogones in the apices of the branches; in

the fact that in many of the branches the cells are no longer than those of the main filament or are shorter; in its secondary branching, and in its branching not being distinctly unilateral.

The genera *Hapalosiphon*, *Stigonema*, and *Fischerella* are in many respects alike. In the table on the next page their chief characteristics may be compared at a glance. This table gives a little more latitude than some of the best writers allow; O. Kirchner, for instance, does not admit secondary branching in *Hapalosiphon*. Different accounts of the same species in these genera often vary greatly, but a safe ground has been taken in compiling this table.

It will be seen from this summary that these genera are separated from each other by no clear and definite lines. There are several intergrading forms. The characters that are used to distinguish them are as follows:

Hapalosiphon grows free-floating in fresh water.

Fischerella grows in moist earth, on wet stones, and in hot springs.

Stigonema grows free-floating or on wet stones.

Stigonema is usually much wider than either of the other two; is often composed of several rows of cells, and the cells are usually more rounded than in *Hapalosiphon*.

The branches in *Fischerella* come from a creeping main filament and are unilateral. In *Hapalosiphon*, the branching is unilateral and secondary branching is relatively uncommon, while in *Stigonema* the filaments may be repeatedly branched in every direction. In contrast to *Stigonema*, *Hapalosiphon* has the cells of its branches longer than those of the main filament. A brown or black sheath is characteristic of *Stigonema*, but some of its species have a colorless sheath.

The question now is whether these differences are sufficient to separate three genera, especially after considering the characteristics of the two new plants above described. Bornet and Flahault in "Revision des Nostocacées Hétérocystées," 1887, class those forms now known as *Stigonema* under the subgenus *Sirosiphon*, and treat *Fischerella* as another subgenus of the genus *Stigonema*. Later, 1895, Gomont proposed raising *Fischerella* to the rank of genus, basing his proposal "sur la différenciation très marquée des filaments primaires rampants, relativement aux filaments dressés." West, in "British Fresh Water Algae" considers *Fischerella* to belong with *Hapalosiphon*.

Thus, there has already been considerable difference of opinion as to the relationship of these forms. *Stigonema anomalum* shows simi-

	Reproduction	Width of Filament	Length of Hormogones	Number of rows in Filaments	Style of Branching	Sheath	Cells	Heterocysts
Hapalosiphon	Hormogones and spores; chiefly spores	4 to 24 microns	All lengths	One, rarely two	Chiefly unilateral; secondary branching relatively uncommon	Smooth; colorless	Generally rectangular	Intercalary
Fischerella	Hormogones and spores	6 to 13 microns	Very long	One or two	Unilateral; branches from creeping main filament; no secondary branching	Smooth; colorless	Rectangular, or rounded	Intercalary
Stigonema	Hormogones and spores; chiefly hormogones	7 to 90 microns	All lengths	One, two, or more.	In every direction; frequent secondary branching	Smooth or rough; colorless or brown to black	Rounded; less commonly rectangular	Intercalary or lateral

larity to *Hapalosiphon* in that its cells are commonly nearly rectangular; its sheath is smooth; there is a distinct tendency to unilateral branching; and the first cell of a branch usually projects into the main filament as in *Hapalosiphon*. *S. medium* shows itself distinctly like a *Stigonema* in its repeated branching; in the fact that the branches do not taper in the least and are often slightly clavate; and in the fact that hormogones are formed in abundance in all the branches. It also differs from the *Hapalosiphon* in having the cells of the branches often shorter than those of the main filament. It therefore seems necessary to put these three genera together as subgenera of the genus *Stigonema*, and let *Sirosiphon* be the subgenus comprising those forms now known as *Stigonema*. Thus:—

Genus, Stigonema Agardh.

1. Subgenus Sirosiphon Bornet & Flahault.
2. Subgenus **Hapalosiphon** (Nägeli) subg. nov.
   Nägeli ex Bornet & Flahault, Revision des Nostocacées Heterocystées, part 3, p. 54, 1887, as genus.
3. Subgenus Fischerella Bornet & Flahault.

Acknowledgement is due to Mr. Frank S. Collins for loan of books and assistance by suggestion, to Mr. Charles Bullard for loan of herbarium specimens, and to Dr. F. D. Lambert for material and assistance.

Bibliography.

Bornet and Flahault: Revision des Nostocacées Heterocystées; part 3: 1887.
West, G. S.: British Freshwater Algae, Cambridge, 1904.
Kirchner, O., in Engler and Prantl, Naturliche Pflanzenfamilien, Teil 1. Abt. 1a, 1898.
Tilden: Minnesota Algae, vol. I, 1910.
Wolle: Fresh Water Algae of the United States, 1887.
Wood: A contribution to the history of the Fresh Water Algae of N. A., 1872.
Forti: Sylloge Myxophycearum, 1907.

Explanation of Plate 105.

Magnification: Figures 1 to 8, × 280; 9 to 12, × 420.

Figures 1 to 8 inclusive are *Stigonema anomalum*.

Fig. 1. The beginning of a branch.
" 2. Young branch and intercalary heterocyst.
" 3. Primary filament with short hormogone-bearing branches, unilaterally arranged. Lateral heterocysts and vacuolated cells.
" 4. Escaping hormogones.
" 5. Escaping hormogone.
" 6. A branch after escape of hormogone.
" 7. Hormogones.
" 8. Formalin material showing connections between cells.

Figures 9 to 12 inclusive are *Stigonema medium*.

Fig. 9. Shows that branching is not unilateral; young branch; division of a cell; two cells side by side.
" 10. Hormogone-bearing branch, and heterocyst.
" 11. Escaping hormogone.
" 12. Thickened portion of sheath at end of hormogone.

Agricultural College, Amherst, Massachusetts.

SIX WEEKS' BOTANIZING IN VERMONT,— II.

ADDITIONAL NOTES ON PLANTS NEAR BURLINGTON.[1]

Sidney F. Blake.

Osmunda cinnamomea L. f. incisa (Huntington) Gilbert. Alt. 3670, Mt. Mansfield, 11 August (*Blake* 2733).

Picea mariana (Mill.) BSP. f. **semiprostrata** (Peck) n. comb. *P. brevifolia* var. *semiprostrata* Peck, Spruces of the Adirondacks 12, 13 (1897).— In moss, alt. 3900 ft., Mt. Mansfield, Underhill, 11 Aug. 1911 (*Blake* 2745).— A creeping alpine form of the Black Spruce, eight or ten decimeters long or more, with short crowded branches mostly fascicled toward the erect apex of the stem, and tiny four-grooved glaucous needles 3–6 mm. long.

Sagittaria arifolia Nutt. Mud flats of Lake Champlain, North Ferrisburg, 8 August 1911.

Cyperus strigosus L. f. **capitatus** (Boeckl.) n. comb. *C. strigosus* var. *capitatus* Boeckl. Linnaea xxxvi. 347 (1869–1870). *C. capitatus* Smyth, Trans. Kansas Acad. Sci. xvi. 163 (1899).— Pasture, Cobble Hill, Milton, growing with the typical form.— A form not very well marked, often — as in Boeckler's original specimens — small and seeming only a dwarfed state of the typical plant, but occasionally larger and more definitely distinguished.

Scirpus atrocinctus Fernald f. brachypodus (Fernald) Blake. Pasture, alt. 1400, Mt. Mansfield, Underhill.

S. heterochaetus Chase. Meadow, North Ferrisburg.

[1] The notes of this list relate to plants collected on several short excursions from Essex Junction, outside the limits of the Burlington Region as defined by Mrs. Flynn.

Juncus effusus L. var. solutus Fernald & Wiegand. Beside wood road, alt. 1260 ft., Mt. Mansfield, Underhill, 11 August (*Blake* 2757).

J. Torreyi Coville. Roadside, alt. 360, Charlotte, 8 August (2612). Previously collected in Vermont by Kirk, in July 1908, in sandy ditch by railroad at Rutland.

Salix sericea Marsh. Mud flats of Hinesburg Pond, Hinesburg, alt. 684, 7 August (2594).

Betula alba L. Woods, alt. 3500, and rocky summit, alt. 3860, Mt. Mansfield, Underhill.

B. alba L. var. cordifolia (Regel) Fernald. Mt. Mansfield, Underhill, alt. 3670.

Parietaria pensylvanica Muhl. Dry soil, alt. 600, Mt. Philo, Charlotte.

Polygonum amphibium L. f. Hartwrightii (Gray) Blake. Mud flats of Hinesburg Pond, Hinesburg (2596).

Epilobium densum Raf. Damp pasture, alt. 465, Milton.

E. molle Torr. With the last.

Sanicula trifoliata Bicknell. Woods, alt. 810, Mt. Philo, Charlotte.

Vaccinium uliginosum L. and var. pubescens Lange. Like so many other species of our blueberries, *Vaccinium uliginosum* possesses both a glabrous and a pubescent form, which have been long distinguished in Europe but not hitherto separated in this country. The plant with the leaves more or less puberulent, sometimes extremely so, beneath, which is V. uliginosum var. pubescens Lange, Consp. Fl. Groenl. 90 (1880), seems with us to be about as common and well distributed as the typical glabrous *V. uliginosum* L. I have seen Vermont material of *V. uliginosum* from Mt. Mansfield, and of var. *pubescens* from Mt. Mansfield, Jay Peak, and Johnson.

Pycnanthemum virginianum (L.) Durand & Jackson. Dry soil, Mt. Philo, Charlotte, alt. 595.

Verbascum Blattaria L. Pasture, North Ferrisburg.

Antennaria Brainerdii Fernald. Dry soil, Mt. Philo, alt. 500.

Eupatorium purpureum L. Woods, alt. 680, Mt. Philo, 8 August 1911 (*Blake* 2611). Second record for the state.

Stoughton, Massachusetts.

A new Station for Scirpus Longii.— Since the publication of the very distinct *Scirpus Longii*[1] in 1911, many students of our flora have sought the plant at favorable localities between the two regions from which it was described,— the Pine Barrens of New Jersey and the Charles River valley in eastern Massachusetts — but so far as the writer has learned quite without success. In the original discussion of the plant it was pointed out that the occurrence of species characteristic of the Pine Barrens is not unprecedented in the valleys of the Charles and the adjacent small rivers, the Neponset, Concord, Mystic, &c., where numerous plants of Coastal Plain distribution occur at isolated stations. It is therefore of at least local interest to record *Scirpus Longii* from the Concord River. In organizing the herbarium of the late Edward S. Hoar, recently presented to the New England Botanical Club, many plants of unusual local interest have been found, collected either by Mr. Hoar or by his intimate friend, Thoreau. Among the sedges are two fine sheets of *Scirpus Longii* collected by Thoreau in 1859 and bearing the original penciled labels "*Scirpus sylvaticus?* [later scratched and marked "*Eriophorum*"] Grt. meadows, May 28" and "*Scirpus Eriophorum.* Great Meadows, July 17, '59." The earlier plant is beginning to flower; the later is fully developed, with good fruit and lingering anthers.[2]— M. L. Fernald, Gray Herbarium.

The Josselyn Botanical Society of Maine met at Thomaston, Tuesday, August 12, 1913 for its Nineteenth Annual Meeting and Field Day. The sessions and field work continued through Wednesday, Thursday and Friday. Collections were made over quite a large area including portions of Thomaston, Rockland, Rockport, Camden, South Thomaston, St. George, Warren and Islesboro. Twenty-five members and guests were in attendance, and by dividing the company into small parties for visiting different localities a large field was covered. Excursions were made to Mt. Megunticook in Camden, to the Lily Pond in Rockport, a large bog in Rockland, to the "Indian Garden" in Warren and to Isleboro and Spruce Head and Elwell Point.

[1] Fernald, Rhodora, xiii. 6 (1911).

[2] Since this note went into type, the writer, while crossing the West Cambridge marshes, on July 8, 1913, came upon a large colony of *Scirpus Longii*, thus demonstrating its presence in the valley of the Mystic River.

The Lily Pond, its shores and the abandoned lime quarries near yielded *Linum catharticum*, *Daphne mezerium* in fine fruit, several species of *Potamogeton* of unusual occurrence which have not yet been fully determined, *Scirpus occidentalis*, and *Dirca palustris*.

Among the species collected at Isleboro were *Dianthus plumarius*, a large form of *Deschampsia caespitosa*, *Galium labradoricum* and *Fragaria virginiana*, var. *terrae-novae*.

The "Indian Garden" is a rich alluvial tract on the St. Georges river, where, according to tradition, the medicine man had his garden of herbs. In close proximity to this is a field from which innumerable arrow-heads and other relics have been taken. In this vicinity were seen several trees of *Quercus macrocarpa*, *Potamogeton americanus*, var. *novaeboracensis* (the first station east of the Connecticut), and a profusion of river-bottom plants growing in great luxuriance.

The party visiting South Thomaston and Elwell Point collected *Euphrasia Randii*, var. *Farlowii*, *Suaeda linearis*, *Distichlis spicata*, *Sparganium americanum* and *Glyceria laxa*.

Collections made during the four days added more than four hundred species from this locality to the Herbarium of the Portland Society of Natural History.

On Thursday evening, August 14, Prof. Fernald gave a highly interesting and important talk on "The Coastal Plain Element in the Maine Flora."

The other evenings were devoted to business meetings and to informal discussion of collections.— D. W. Fellows, Secretary.

The Nineteenth Annual Field Meeting of the Vermont Botanical Club was held at Townshend, Vermont, July 1 and 2, in conjunction with the Vermont Bird Club, with an attendance of thirty.

The first day was spent by some in climbing Ball Mountain where nothing of particular botanical interest was seen. Others explored the banks of the West River where *Sanguisorba canadensis*, *Habenaria flava*, *Prunus pumila* and various interesting carices were found.

The second day was spent in a trip to the top of Newfane Hill where Prof. A. J. Grout entertained the Clubs, at his log cabin and showed them stations for some rare mosses.

July 3 a party which varied in number from thirteen to nineteen

went on a camping trip into the wilds of Stratton about fifteen miles from Townshend. The headquarters were at Grout's Mill the place where six of the members on July 4, 1895, formed a temporary organization which later resulted in the Club, which has grown to a membership of over two hundred.

During the week following, trips were made to Stratton Mountain and to several ponds and old lumbering operations within tramping distance. *Lobelia Dortmanna, Sisyrinchium atlanticum, Habenaria fimbriata, Carex limosa, C. Michauxiana, C. lenticularis* and other interesting plants were secured.

Particular attention was paid to the blackberries. Mr. W. H. Blanchard's study of the Vermont members of this genus, was made chiefly in Windham Co., and several members of the Club, aware of this fact, visited his stations at Townshend, which were indicated with exactness in a letter to Mr. Wheeler, and his eight new species for this region recognized in the new Gray Manual were all seen. Later the study was carried on with much enthusiasm by the party at Stratton and all of his species for that region were found.

The weather was all that could be hoped for and a very profitable time was spent.— Mrs. Nellie F. Flynn, Burlington, Vermont.

A correction concerning Dicksonia punctilobula, forma cristata. — It is necessary to correct the report of the collection of *Dicksonia punctilobula* (Michx.) Gray, forma *cristata* (Maxon) Clute, which was made in Rhodora, XV, 44 (1913). The report was made from a mistaken determination, and the specimen proves to be merely a forked state of the typical form. — Martha Louise Loomis, Sherborn, Massachusetts.

Vol. 15, no. 178, including pages 173 to 188, was issued 17 October, 1913.



Figs. 1–8. Stigonema anomalum Blanchard, sp. nov.
Figs. 9–12. Stigonema medium Blanchard, sp. nov.

Just Published

COULTER'S PLANT LIFE AND PLANT USES

An Elementary Textbook: Foundational to Agriculture, Domestic Science, or College Botany

PRICE, $1.20

A high school book about the fundamentals of plant life and about the relation between plants and man. It aims to present only those large facts about plants which form the present basis of the science of botany, and which are essential to every high school botanist. In so far as is possible, plant life is presented in terms of its largest relation to life, but the treatment has in view preparation for life in general, and not preparation for any particular kind of calling. The subject matter has been presented in an interesting manner and in as simple a way as possible; scientific words, where first used, are interpreted by simple English equivalents. The illustrations are abundant and excellent.

AMERICAN BOOK COMPANY

NEW YORK CINCINNATI CHICAGO

HOME GROWN LILIES.

Fresh from Beds. Wild Flowers, Hardy plants of all kinds.
Send for catalogue.

F. H. HORSFORD, Charlotte, Vt.

BOTANY LABELS 85c. PER 1000

POSTPAID

Also, printed envelopes, letterheads and statements
$1.00 per 1000 plus stock

For samples write to D. LEWIS DUTTON, Brandon, Vermont

CAMBRIDGE BOTANICAL SUPPLY COMPANY,

WAVERLEY, MASS.

BOTANICAL SUPPLIES OF EVERY DESCRIPTION.
Samples of Driers and Mounting Papers sent on request.
ASK FOR NEW CATALOG, NO. 91.

Standard Material. **Immediate Shipment.**